



Efficient reduction of CO₂ using a novel carbonic anhydrase producing *Corynebacterium flavescens*

Tanvi Sharma, Ashok Kumar[†]

Department of Biotechnology and Bioinformatics, Jaypee University of Information Technology, Waknaghat, Solan, India

Abstract

Emission of greenhouse gases into the atmosphere by human activities leads to global warming. To reduce the level of CO₂ the bio-catalytic properties of microbial carbonic anhydrase (CA) can be exploited. The present study aimed to isolate CA producing bacteria from cow saliva. After thorough screening for CA activity in the bacterial cultures, ten isolates were selected. Out of ten bacterial isolates, T5 isolate showed the highest CA activity (83.92 U/mL) and the isolate was identified as *Corynebacterium flavescens* using 16s rRNA analysis. Various production parameters for the optimum production of CA were optimized. During the optimization, incubation temperature 40°C, agitation speed 120 rpm, and inoculum volume 4% v/v was found to be optimum for CA production. The optimum reaction pH, reaction time and temperature were 7, 10 min and 35°C, respectively. The crude enzyme was tested for the conversion of CO₂ into the calcium carbonate (CaCO₃) under controlled conditions. The CO₂ conversion efficacy of crude CA was observed to be ~45 mg CaCO₃/mg protein. The synthesized CaCO₃ was analyzed using scanning electron microscopy and X-ray diffraction techniques for particle size, morphology and elemental structure. Calcite precipitation by bacterial CA makes it a potential candidate to be effectively employed in biomimetic CO₂ sequestration.

Keywords: Carbonic anhydrase, CO₂ sequestration, *Corynebacterium flavescens*, SEM



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>)

which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received April 21, 2020 Accepted July 08, 2020

[†] Corresponding Author

E-mail: ashok.nadda09@gmail.com

Tel: +01792-239351-353 Fax: +01792-239351

Orchid: 0000-0001-9192-0774

1. Introduction

Global warming is one of the serious issues due to excessive emission of carbon dioxide (CO_2) into the atmosphere. To reduce CO_2 emission, numerous government and non-governmental organizations endorse to switch from fossil fuel power to clean energy sources to protect the atmosphere from the terrible effects of global warming [1-3]. Furthermore, worldwide research is going on for finding a potent and eco-friendly method that directly converts industrial emitted CO_2 into a useful product. Some of the algae, bacteria, and cyanobacteria play a vital role in alleviating the increasing level of CO_2 using their carboxylating enzyme [4, 5]. Recently, these carboxylating enzymes are getting attention due to its role in CO_2 sequestration [1, 6]. However, algae have good potential for CO_2 fixation, but it has a few disadvantages such as light requirement [7]. Hence, it is advantageous to isolate a CO_2 fixing bacteria or enzyme which does not require continuous light for its growth and activity [8].

CA is one of the fastest known biocatalysts and involved in the CO_2 sequestration. CA contains zinc in its active center and converts CO_2 to bicarbonates, which can be further converted to CaCO_3 in the presence of calcium ion. The application of CA in the conversion of CO_2 from flue gas into thermodynamically stable, environmentally safe calcium carbonate offers several advantages [9]. First of all, the process is extremely fast and take place near ambient condition. CaCO_3 formed during CO_2 conversion can be utilized in the preparation of white pigment, cement, antacids and others [10]. Thus, a new biomimetic approach using CA has been found to be viable for fixing a huge quantity of CO_2 into CaCO_3 . CA catalyzes several other hydrolytic reactions too, including hydration of urea, carboxylic acid, halides, and hydrolyzable substrates [11, 12]. The CA catalyze the hydration of CO_2 in two steps ping-pong mechanism. Its

1 active site has divalent metal ions generally zinc in a tetrahedral confirmation, containing three
2 amino acid as ligands and hydroxide ions coordinating the metal [13]. CA is present in
3 eukaryotes and prokaryotes, where it takes part in various physiological functions such as acid-
4 base balance, hemostasis, respiration, and photosynthetic CO₂ fixation [14]. Although CA is
5 present ubiquitously but microbial CA has received more attention due to ease of their
6 production and applications. Till now, CA has been classified into five (α , β , γ , δ and ϵ) different
7 classes [15]. The first studied, class is the α -CA as it plays a crucial role in human pathology and
8 drug targeting. However, the other classes take part in the conversion of gaseous CO₂ [16]. The β
9 class exists as a dimer, tetramer and hexamer and mostly present in algae, bacteria, and archaea.
10 The γ class is mostly found in archaea and exists as a trimer [17]. This enzyme is either present
11 extracellularly or inside (intracellular) the cytoplasm in the bacteria [18].

12 Recent studies suggest that CA is widespread in bacteria and archaea domain, indicates
13 that this enzyme has a more extensive and vital role in prokaryotic physiology [19].CA
14 producing bacteria were isolated from various sources such as deep sea-water, alkaline soil,
15 mangroves soil, and seashore water [20]. Moreover, the optimization of various parameters is
16 essential to enhance the total yield, maximal activity of the enzyme and to reduce the process
17 cost. The high enzyme titer has been attained by changing the ratio of various media ingredients
18 that are influencing the growth and enzyme production in bacteria. The optimization is greatly
19 influenced by various physicochemical factors and nutritional such as temperature, agitation rate,
20 pH, media components, inoculum volume, *etc.* Thus, optimizing the best conditions for
21 maximum enzyme production is still vital and essential in biocatalytic transformations.

1 In the present study, CA producing bacteria was isolated from cow saliva and the
2 parameters were optimized for the production of CA in bulk amount. The application of crude
3 CA has been studied for the conversion of CO₂ into calcium carbonates (Fig. 1). As per our
4 knowledge, this is the first report on the isolation of CA producing bacteria from cow saliva. In
5 the current global warming scenario, this work might be helpful in CO₂ conversion and
6 environmental amelioration. Here, we demonstrate that isolated bacteria are a green biocatalyst
7 for the conversion of CO₂ into calcium carbonates.

8

9 **2. Methods**

10 **2.1. Isolation of a Bacterial Strain**

11 CA is a ubiquitous enzyme present in all plants, mammalian tissues, algae, and bacteria [19]. In
12 the present study, the sample was collected from six years old cow saliva from District Mandi,
13 Himachal Pradesh, India with the help of sterilized bud (Fig. 2). The saliva was serially diluted
14 and then 100 µL of culture was transferred onto nutrient agar having 3mM *p*-nitrophenyl acetate
15 (*p*-NPA), incubated at 30°C for 48 h. The appearance of the intense yellow colour colonies
16 indicated the production of carbonic anhydrase [21]. The bacterial isolates, which utilized *p*-NPA
17 on agar plates, were further screened for CA activity in nutrient broth. For CA activity assay, the
18 seed culture was prepared aseptically in 50 mL nutrient broth containing a loopful culture of T5
19 isolate and incubated at 30°C for 24 h. For the production of CA, the nutrient broth was used as
20 production media and the seed culture 2% v/v was used as inoculum for 50 mL production
21 medium. The inoculated production medium was incubated at 30°C for 36 h at 120 rpm. The
22 broth was centrifuged at 10,000 rpm for 10 min after 36 h of incubation. Then pellet was

1 suspended in phosphate buffer (pH 7.0) and homogenized well. The suspension was sonicated
2 using 20 kHz frequency for 5 min at 4°C and centrifuged at 10,000 rpm for 20 min. The
3 supernatant was used as crude enzyme extract.

4 5 **2.2. Enzyme Assay**

6 CA activity was assayed in the culture broth using a previously reported method by measuring
7 the micromole of *p*-nitrophenol released from *p*-NPA [22]. To 825 µL of Phosphate buffer
8 (50mM, pH 7.5), 175 µL of the substrate stock solution (*p*-NPA, 10 mM in isopropanol) was
9 added. Then, the reaction mixture was incubated in a water bath at 37°C for 5 min. Subsequently
10 to initiate the reaction 25 µL of an enzyme was added. The reaction mixture was again incubated
11 37°C in a water bath for 5 min [23]. The amount of *p*-nitrophenol released was measured at 410
12 nm after 5-min incubation using a microplate reader (Thermo-scientific, Multiskan™ FC
13 Microplate Photometer) [24]. Based on CA activity, a potent isolate T5 was selected for all
14 subsequent studies. The colonies of T5 isolate was subjected to gram's staining and observed at
15 100 X magnification under a compound microscope. The biochemical characterization of T5
16 isolate was conducted according to Bergey's Manual of Systemic Bacteriology [25].

17 18 **2.3. Identification of Bacterial Strain**

19 The bacterial strain was identified by 16S rRNA sequencing by Bioreserve Biotechnologies Pvt.
20 Ltd., Hyderabad, India [26]. The nucleotide sequences obtained from 16s rRNA sequencing were
21 subjected to a homology search using BLAST and aligned using a MEGA X software. Then, the
22 phylogenetic tree was made using the neighbour-joining method.

1 **2.4. Optimization of Production Parameter for CA Producing Bacteria**

2 Luria broth, nutrient broth, peptone broth, minimal salt media, muller Hinton broth, and basal
3 salt media were used for enzyme production. Luria broth, nutrient broth, and muller hinton broth
4 were procured from Hi Media, Mumbai. The composition of other media used for enzyme
5 production was peptone broth (g/L): beef extract 3.0, glucose 1.0, NaCl 5.0, peptone 5.0, CaCO₃
6 6.0; minimal salt media (g/L): KH₂PO₄ 6.8, MgSO₄ 0.2, Na₂HPO₄, 7.8, ZnCl₂ 0.02,
7 ZnSO₄.7H₂O 0.05, NaNO₃, 0.085; and basal slat media (g/L): sucrose 5.0, MgSO₄.7H₂O 0.5,
8 Na₂HPO₄ 2.0, FeCl₃·6H₂O 0.005, CaCO₃ 0.1. The effect of temperature on the CA production
9 was studied by incubating production culture at various temperatures ranging from 20 to 60°C.
10 To determine the effect of inoculum volume (% v/v) on CA production, production media was
11 inoculated with a (2-12 %v/v) culture of T5 isolate and incubated at 40°C. The optimum
12 agitation rate for CA production was determined by incubating the production media at 40°C at
13 varying agitation speeds (80, 100, 120, 140, 160, and 180 rpm). The enzyme activity was
14 determined for each incubated production media sample using standard activity assay [23].
15

16 **2.5. Optimization of Reaction Parameter for CA Producing Bacteria**

17 For optimizing reaction conditions of CA effect of reaction time (2-18 min), different buffer
18 systems (citrate buffer, potassium phosphate buffer, and Tris–HCl buffer, 50 mM), reaction
19 temperature (25-60°C), different metal ions (Na⁺, K⁺, Ca²⁺, Zn²⁺, Mg²⁺, Fe³⁺, and Al³⁺) and
20 organic solvent (ethanol, isopropanol, ethanediol, *n*-butanol, propandiol, and, acetonitrile) was
21 studied on CA activity.
22

1 **2.6. Enzymatic Conversion of Carbon Dioxide into Calcite**

2 The potential of CA for CO₂ conversion was determined as described by Giri et al. [27], with
3 minor modifications. The precipitation of CO₂ into calcium carbonate (CaCO₃) was carried out
4 in a 50 mL total reaction mixture containing 2 mL Tris buffer (1M, pH 8), 23 mL of 2 (%v/v)
5 calcium chloride solution, 23 mL of CO₂ saturated water and 2 mL (0.5 mg/mL) of an enzyme in
6 phosphate buffer (50 mM, pH 7). The reaction was performed for 10 min at room temperature.
7 Bovine serum albumin (BSA) was used as a control. The precipitate formed after 10 min was
8 recovered by centrifugation. The sediment of precipitates was lyophilized overnight to obtain the
9 dry powder. Then, CaCO₃ precipitates were weighed to determine the amount of carbonate
10 deposited during the enzymatic reaction.

11 12 **2.7. Instrumental Analysis of CaCO₃**

13 The CaCO₃ precipitates were lyophilized to get dry powder of CaCO₃. The CaCO₃ precipitates
14 were scanned in the range of 400-4,000 cm⁻¹. Compositions of the precipitated solid crystals
15 were examined by using X-ray diffraction (XRD). To determine crystal morphology Scanning
16 Electron Microscopy (SEM) was performed by using JSM7401F (JEOL).

17 18 **3. Results and Discussion**

19 **3.1. Isolation of CA Producing Bacteria**

20 The ten bacterial isolates from a cow saliva sample, which efficiently utilized *p*-NPA on agar
21 plates, were selected for secondary screening and cultivated in nutrient broth to detect CA
22 activity. *p*-NPA plates assay is specific and based on the formation of yellow-colored (*p*-NP)

1 product formed around the colonies. Among these bacterial isolate T5, showed maximum CA
2 activity, and selected for further work (Fig. 3 (a)). The T5 isolate was rod-shaped and gram-
3 positive bacilli (Fig. 3 (b)). And the results of the biochemical analysis show Voges-Proskauer
4 positive, nitrate reduction positive, glucose and sucrose positive. The presence of CA in
5 mammalian saliva is reported several decades ago. Yoshimura et al. [28] suggested that the role
6 of CA in mammalian saliva is to regulate saliva pH. Furthermore, the screening of
7 microorganisms from cow saliva for CA production is the rapid and cost-effective method. Jaya
8 et al. [29] reported an isolate *Bacillus safensis* from the water sample that produced CA.
9 Similarly, Kupriyanova et al. [30] reported that the presence of CA in *Microcoleus*
10 *cathonoplastes*, isolated from the soda lakes in Russia. Various pathogenic bacteria such as
11 *Yersinia pseudotuberculosis* and *Listeria monocytogenes* isolated from soil and water were
12 reported to produce CA [31]. Microorganisms need CO₂ and bicarbonate in their metabolic
13 activities and CA plays a vital role to meet these demands.

14

15 **3.2. Identification of Strain**

16 The genomic DNA was isolated from bacterial isolate T5 and its DNA analysis was done by
17 agarose gel electrophoresis (Fig. 4 (a)). The 16s rRNA was amplified from genomic DNA using
18 universal primers. Then bacterial strain was identified as *Corynebacterium flavescens* by 16S
19 rRNA sequencing based on phylogenetic analysis and nucleotide homology. The phylogenetic
20 tree was made by using a neighbour tree joining method showing maximum sequence homology
21 of 99.42% and 99.27% with *Corynebacterium flavescens* HBUM07012 and *Corynebacterium*

1 *flavescens* EHFS1S12Hc, respectively (Fig. 4 (b)). The gene accession number of the organism
2 was MN982752 deposited in NCBI.

3

4 **3.3. Optimization of Production Conditions for CA**

5 optimization of production and reaction parameters is widely known to rise the production
6 significantly. In the present study, out of six media, the maximum CA activity (84.99 U/mL) was
7 observed in nutrient broth media (Fig. 5 (a)). In contrast, to the present findings, Sharma et al.
8 [23] reported maximum CA production from *Pseudomonas fragi* in peptone broth. Also, in a
9 previous study, the nutrient broth was reported as the best medium for the production of CA
10 from *Nocardiopsis lucentensis* [20]. Thus, the nutrient broth media was selected for further
11 studies. The temperature of incubation significantly affects the production of microbial enzymes
12 and their activity. The maximum production of CA with residual activity 85.59 U/mL was
13 observed at 40°C. Furthermore, a decrease in the CA activity was noticed with a further increase
14 in temperature (Fig. 5 (b)) [24]. The decrease in activity might be attributed to the denaturation
15 of protein at a higher temperature. Similarly, Sharma et al. [23], reported the optimum CA
16 production was observed at 40°C. In a recent study, *Bacillus safenis* was observed to show
17 optimum activity at 40°C [29]. In another study, CA from *Citrobacter freundii* was found to be
18 active at 37°C [27]. However, the optimum temperature for CA production from
19 *Methanobacterium thermoautotrophicum* was found to be 75°C [22]. These results showed that
20 CA production has its own temperature optima and is species-specific which favours maximum
21 CA production.

1 The volume of inoculum also affected the production and activity of enzymes from the
2 isolated bacterial strain. An increase in CA production was observed with an increase in an
3 inoculum volume up to 4% v/v then it started decreasing (Fig. 5 (c)). Another study showed that
4 1.5% inoculum size was optimum for CA production from *Aeribacillus pallidus* [32]. Usually
5 increased inoculum volume improves the growth of bacteria up to a certain level and after that
6 bacterial growth starts decreasing due to nutritional limitation. Furthermore, lower inoculum
7 volume has a low number of bacterial cells in medium and this requires a longer time to grow.
8 This might lead to the accumulation of toxic compounds which ultimately decreased enzyme
9 production [33]. However, some studies reveal that an increase in inoculum size results in more
10 enzyme production and *vice versa*. The maximum CA activity was observed at an agitation
11 speed of 120 rpm *i.e.*, 86.50 U/mL (Fig. 5 (d)). The CA activity was increased by increasing the
12 agitation rate up to 120 rpm. But the high agitation rate also resulted in decreased activity due to
13 mechanical disruption of proteins. In a previous study, carried out by Zhang et al. [34], showed
14 that 150 rpm most appropriate agitation rate for the production of CA whereas CA from
15 *Aeribacillus pallidus* gave maximum activity at 200 rpm [32]. The key role of agitation in the
16 fermentation broth is to ensure adequate mixing of oxygen that eventually becomes available for
17 the growth of bacteria in the dissolved form of oxygen. Also, microbial enzyme production is
18 observed to be good under continuous shaking conditions as compared to static conditions [35].

19

20 **3.4. Optimization of Reaction Parameters for CA**

21 The effect of pH on CA activity was determined and optimum pH for maximum CA activity was
22 7 in phosphate buffer (Fig. 6 (a)). A decrease in activity at high and low pH generally results in

1 loss of enzyme activity due to disruption of three-dimensional structure and alteration in amino
2 acid residues present in the active center of an enzyme. In another study, CA from *Bacillus sp.*
3 gave maximum activity at pH 8 whereas CA from *Lactobacillus* showed optimum activity at pH
4 6 [36, 37]. However, CA from *Helicobacter pylori* showed optimum activity in an acidic
5 environment and high acid tolerance [38]. The effect of reaction temperature on the CA activity
6 was studied. The result presented in (Fig. 6 (b)) showed that the optimum temperature for CA
7 activity was 35°C after that decline in CA activity was observed. In the previous study, the
8 optimum temperature for purified CA was found to be 60°C [36]. Mostly, CA isolated from
9 human erythrocyte and bovine shows CA activity in the temperature range of 35-40°C and in the
10 pH range of 6.5-7.5 [39]. The optimum reaction time for CA activity was observed after 10 min
11 of incubation (88.55 U/mL) (Fig. 6 (c)). After 10 min decrease in enzyme activity was observed
12 which may be due to the inhibition of product and denaturation of the enzyme. All these facts
13 indicate the functional diversity of CA and the capability of this enzyme to perform diverse roles
14 in the organisms living in extreme environments.

15 Out of various organic solvent used, all of them decrease CA activity (Fig. 6 (d)). This
16 can be credited to conformational changes in enzyme active sites that are responsible for the
17 decrease in enzyme activity [40]. Furthermore, the effect of metal ion on CA activity at
18 concentrations 1mM and 5mM was studied (Table 1). The presence of metal ion in the reaction
19 medium either inhibit or promote the enzymatic reaction. CA activity was found to be increased
20 in presence of Zn^{2+} and Fe^{3+} ion this might be due to reason these metal ions are present in the
21 active site of CA, they could have a stabilization effect thus enhancing the CA activity. In
22 contrast, CA activity was not affected by the presence of Ca^{2+} ion indicates that enzyme could

1 perform in the presence of calcium ion. Whereas, CA activity was inhibited by the presence of
2 Na^{2+} , K^+ , Mg^{2+} , and Al^{3+} ions. In a previous study, CA activity from *Pseudomonas fragi* was
3 significantly enhanced by Zn^{2+} , Fe^{3+} and Cd^{2+} , Na^{2+} , K^+ , Mg^{2+} , and Ca^{2+} had no effect, whereas
4 Hg^{2+} and Pb^{2+} had an inhibitory effect [23]. In contrast, the activity of CA from *Dicentrarchus*
5 *labrax* was inhibited by Zn^{2+} , Hg^{2+} , Al^{3+} , Cu^{2+} , and CO^{2+} [41].

7 **3.5. Enzymatic Conversion of Carbon Dioxide Using Crude Enzyme**

8 Finally, the application of CA to convert CO_2 into CaCO_3 in the presence of calcium ion was
9 examined. Test reaction containing crude CA showed a significant conversion of CO_2 as
10 compared to that of control. BSA was used as a control in the experiment which showed no
11 precipitation of CO_2 . The CO_2 conversion efficacy of CA was studied by calculating the amount
12 of CaCO_3 synthesized. The crude CA resulted in the formation of 45 mg CaCO_3 /mg of protein.
13 The CO_2 conversion efficiency of the enzyme produced from *Corynebacterium flavescens* is
14 much higher as compared to CA from *Bacillus pumilis* (33.06 mg CaCO_3 /mg of protein) and
15 *Pseudomonas fragi* (27.33 mg CaCO_3 /mg of protein) [42, 23].

16 CaCO_3 is commonly used in various industries such as fillers for paints, papers, plastics,
17 as well as food and pharmaceutical applications [43]. Therefore, it confirmed that the bacterial
18 isolate is producing potent CA which can be used for the CO_2 conversion in the lab as well as
19 commercial scale.

21 **3.6. Instrumental Analysis of CaCO_3**

1 In the present studies, the crude enzyme precipitates CaCO_3 and exhibited morphological
2 resemblance with CaCO_3 standard taken from Sigma Aldrich. In the SEM analysis, spherical
3 vaterite crystal was formed in control and rhombohedral vaterite and calcite crystals were formed
4 in the presence of an enzyme (Fig. 7 (a) and (b)). This result from SEM analysis conforms to the
5 finding of Jo et al. [44]. The composition of precipitated powder was confirmed by XRD
6 analysis. The two forms of CaCO_3 were vaterite, and calcite, analyzed by XRD patterns.
7 Diffraction peak at 29.38, 34.04, 43.90 corresponds to the calcite crystal phase whereas the peak
8 at 22.54, 27.08 corresponds to the vaterite crystal phase (Fig. 7 (c)). Furthermore, various studies
9 have shown that calcite and vaterite were phase formed in the presence of CA [37, 44].

10

11 **4. Conclusions**

12 Microorganisms are commonly used to produce the enzymes of industrial utility. The isolation of
13 CA from cow saliva is cost-effective as the price of commercially BCA is about $\$3,000 \text{ g}^{-1}$. Thus,
14 there is a price limitation for using BCA in various CO_2 conversion applications. The improved
15 CA production from *Corynebacterium flavescens* was achieved by the optimization of various
16 production and reaction parameters. The high yield of CA was achieved from this process as an
17 indication of the possibility for its large-scale production. The crude enzyme was also found to
18 be effective in the CO_2 conversion experiment and hence its efficacy in CO_2 capture can be
19 exploited at the industrial level too. In future studies, the CA enzyme can be purified and used in
20 an immobilized form for better reaction efficiency. Moreover, directed evolution and rational
21 hybrid design can be used to develop the mutant of this enzyme with better enzyme stability for
22 industrial purposes.

1

2 **Acknowledgment**

3 The financial support from the Jaypee University of Information Technology, Waknaghat to
4 undertake this study is thankfully acknowledged. Further, the authors have no conflict of interest
5 either among themselves or with the parent institution.

6

7 **Author Contributions**

8 T.S. (Ph.D student) performed all the experiments and wrote the manuscript.

9 A.K. (Assistant Professor) designed experiments and revised the manuscript.

10

11 **References**

12 1. Sharma T, Sharma S, Kamyab H, et al. Energizing the CO₂ utilization by chemo-enzymatic
13 approaches and potentiality of carbonic anhydrases: A review. *J. Clean. Prod.* 2020;247:1-12.

14 2. Zhang W, Liu H, Sun C, et al. Capturing CO₂ from ambient air using a polyethyleneimine–
15 silica adsorbent in fluidized beds. *Chem. Eng. Sci.* 2014;116:306-316.

16 3. Mulenga D, Siziya S. Indoor Air Pollution Related Respiratory Ill Health, a Sequel of Biomass
17 Use. *SciMed. J.* 2019;30-37.

18 4. DiMario RJ, Machingura MC, Waldrop GL et al. The many types of carbonic anhydrases in
19 photosynthetic organisms. *Plant Sci.* 2018;268:11–17.

20 5. Kupriyanova EV, Sinetova MA, Cho SM, et al. CO₂-concentrating mechanism in
21 cyanobacterial photosynthesis: organization, physiological role, and evolutionary origin.

22 *Photosynth Res.* 2013;117(1-3):133-46.

- 1 6. Chen F, Jin W, Gao H, et al. Cloning, Expression and Characterization of Two Beta Carbonic
2 Anhydrases from a Newly Isolated CO₂ Fixer, *Serratia marcescens* Wy064. *Indian J. Microbiol.*
3 2018;59(1):64-72.
- 4 7. Rinanti A. Biotechnology carbon capture and storage by microalgae to enhance CO₂ removal
5 efficiency in closed-system photobioreactor. In: Thajuddin N and Dhanasekaran D eds, *Algae-*
6 *Organisms for Imminent Biotechnology*. IntechOpen; 2016. p.134-136. 10.5772/62915.
- 7 8. Hou J, Li X, Kaczmarek MB, et al. Accelerated CO₂ Hydration with Thermostable
8 *Sulfurihydrogenibium azorense* Carbonic Anhydrase-Chitin Binding Domain Fusion Protein
9 Immobilised on Chitin Support. *Int. J. Mol Sci.* 2019;20(6):1494.
- 10 9. Chang R, Kim S, Lee S, et al. Calcium carbonate precipitation for CO₂ storage and utilization:
11 A review of the carbonate crystallization and polymorphism. *Front. Energy Res.* 5 2017;1-17.
- 12 10. Faridi S, Satyanarayana T. Applicability of carbonic anhydrase in mitigating global warming
13 and development of useful products from CO₂. *Clim. Change Environ. Sustain.* 2015;3:77-92.
- 14 11. Bose H, Satyanarayana T. Utility of thermo-alkali-stable gamma-CA from polyextremophilic
15 bacterium *Aeribacillus pallidus* TSHB1 in biomimetic sequestration of CO₂ and as a virtual
16 peroxidase. *Environ. Sci. Pollut. Res. Int.* 2017;24(11):10869-10884.
- 17 12. Kumar A, Sharma T, Mulla SI, et al. Let's Protect Our Earth: Environmental Challenges and
18 Implications. In: Kumar A, Sharma S eds. *Microbes and Enzymes in Soil Health and*
19 *Bioremediation*. Singapore: Springer Singapore; 2019. p. 1-10.
- 20 13. Aggarwal M, Chua TK, Pinard MA, et al. Carbon Dioxide "Trapped" in a β -Carbonic
21 Anhydrase. *Biochemistry* 2015;54(43):6631-6638.


- 1 14. Abo-Ashour MF, Eldehna WM, Nocentini A, et al. Novel synthesized SLC-0111 thiazole
2 and thiadiazole analogues: Determination of their carbonic anhydrase inhibitory activity and
3 molecular modeling studies. *Bioorg. Chem.* 2019;87:794-802.
- 4 15. Alissa SA, Alghulikah HA, Alothman ZA, et al. Phosphoramidates are the first phosphorus-
5 based zinc binding motif to show inhibition of β -class carbonic anhydrases from bacteria, fungi,
6 and protozoa. *J. Enzyme. Inhib. Med. Chem.* 2020;35(1):59-64.
- 7 16. Boone CD, Habibzadegan A, Gill S, et al. Carbonic anhydrases and their biotechnological
8 applications. *Biomolecules* 2013;3(3):553-62.
- 9 17. Prete SD, Nocentini A, Supuran CT. Bacterial α -carbonic anhydrase: a new active class of
10 carbonic anhydrase identified in the genome of the Gram-negative bacterium *Burkholderia*
11 *territorii*. *J. Enzym. Inhib. Med. Ch.* 2020;35(1):1060-1068.
- 12 18. Hu JJ, Wang L, Zhang SP, et al. Optimization of electron donors to improve CO₂ fixation
13 efficiency by a non-photosynthetic microbial community under aerobic condition using statistical
14 experimental design. *Bioresour. Technol.* 2010;101(18):7073-7078.
- 15 19. Smith KS, Ferry JG. Prokaryotic carbonic anhydrases. *FEMS Microbiol. Rev.* 2000;24:335-
16 366.
- 17 20. Anjana G, Dudhagara P, Bhagat C, et al. Biomimetic Sequestration of CO₂ Using Carbonic
18 Anhydrase from Calcite Encrust Forming Marine Actinomycetes. *Sci. Int.* 2015;3:48-57.
- 19 21. Ramanan R, Kannan K, Sivanesan SD, et al. Bio-sequestration of carbon dioxide using
20 carbonic anhydrase enzyme purified from *Citrobacter freundii*. *World. J. Microb. Biot.*
21 2009;25(6):981-987.

- 1 22. Smith KS, Ferry JG. A plant-type (beta-class) carbonic anhydrase in the thermophilic
2 methanoarchaeon *Methanobacterium thermoautotrophicum*. *J. Bacteriol.* 1999;181(20):6247-
3 6253.
- 4 23. Sharma A, Bhattacharya A, Singh S. Purification and characterization of an extracellular
5 carbonic anhydrase from *Pseudomonas fragi*. *Process. Biochem.* 2009;44(11):1293-1297.
- 6 24. Sharma A, Sharma T, Meena KR, et al. High throughput synthesis of ethyl pyruvate by
7 employing superparamagnetic iron nanoparticles-bound esterase. *Process. Biochem.*
8 2018;71:109-117.
- 9 25. Meena K, Sharma A, Kumar R, et al. Two factor at a time approach by response surface
10 methodology to aggrandize the *Bacillus subtilis* KLP2015 surfactin lipopeptide to use as
11 antifungal agent. *J. King Saud Univ. Sci.* 2018;32(1):337-348.
- 12 26. Sharma A, Meena K, Kanwar S. Molecular characterization and bioinformatics studies of a
13 lipase from *Bacillus thermoamylovorans* BHK67. *Int. J. Biol. Macromol.* 2017;107:2131-2140.
- 14 27. Giri A, Banerjee UC, Kumar M, et al. Intracellular carbonic anhydrase from *Citrobacter*
15 *freundii* and its role in bio-sequestration. *Bioresour. Technol.* 2018;267:789-792.
- 16 28. Yoshimura H, Iwasaki H, Nishikawa T, et al. Role of carbonic anhydrase in the bicarbonate
17 excretion from salivary glands and mechanism of ionic excretion. *Jpn. J. Physiol.* 1959;9(1):106-
18 23.
- 19 29. Jaya P, Nathan V, Parvathi A. Characterization of marine bacterial carbonic anhydrase and
20 their CO₂ sequestration abilities based on a soil microcosm. *Prep. Biochem. Biotech.*
21 2019;49:891-899.

- 1 30. Kupriyanova E, Villarejo A, Markelova A, et al. Extracellular carbonic anhydrases of
2 stromatolite forming cyanobacterium *Microcoleus cathonoplastes*. *Microbiol.* 2007;153:1149-
3 1156.
- 4 31. Buzolyova LS, Somov G. Autotrophic assimilation of CO₂ and C1 compounds by pathogenic
5 bacteria. *Biochemistry* 1996;4 (10):1357-1361. 10561561.
- 6 32. Bose H, Satyanarayana T. Suitability of the alkalistable carbonic anhydrase from a
7 polyextremophilic bacterium *Aeribacillus pallidus* TSHB1 in biomimetic carbon sequestration.
8 *Bioproc. Biosyst. Eng.* 2016;39(10):1515-1525.
- 9 33. Shivalee A, Lingappa K, Mahesh D. Influence of bioprocess variables on the production of
10 extracellular chitinase under submerged fermentation by *Streptomyces pratensis* strain KLSL55.
11 *J. Genetic Eng. Biotech.* 2018;16(2):421-426.
- 12 34. Zhang Z, Lian B, Hou W, et al. Optimization of nutritional constituents for carbonic
13 anhydrase production by *Bacillus mucilaginosus* K02. *Afr. J. Biotechnol.* 2011;10:8403-8413.
- 14 35. Fleuri L, Sato H. Production, purification, cloning and application of lytic enzymes. *Quím.*
15 *Nova.* 2005;28:871-879.
- 16 36. Sundaram S, Thakur IS. Induction of calcite precipitation through heightened production of
17 extracellular carbonic anhydrase by CO₂ sequestering bacteria. *Bioresour. Technol.*
18 2018;253:368-371.
- 19 37. Li CX, Jiang XC, Qiu YJ, et al. Identification of a new thermostable and alkali-tolerant α -
20 carbonic anhydrase from *Lactobacillus delbrueckii* as a biocatalyst for CO₂ biomineralization.
21 *Bioresour. Bioprocess.* 2015;2(1):44.


- 1 38. Chirica LC, Elleby B, Lindskog S. Cloning, expression and some properties of a-carbonic
2 anhydrase from *Helicobacter pylori*. *Biochim. Biophys. Acta* 2001;1544:55-63.
- 3 39. Demir N, Demir Y, Coskun F. Purification and characterization of carbonic anhydrase from
4 human erythrocyte plasma membrane. *Turk. J. Med. Sci.* 2001;31:477-482.
- 5 40. Kumar A, Dhar K, Kanwar SS, et al. Lipase catalysis in organic solvents: advantages and
6 applications. *Biol. Proced. Online.* 2016;18:2-2.
- 7 41. Ceyhun SB, Senturk M, Yerlikaya E, et al. Purification and characterization of carbonic
8 anhydrase from the teleost fish *Dicentrarchus labrax* (European Seabass) liver and toxicological
9 effects of metals on enzyme activity. *Environ. Toxicol. Pharmacol.* 2011; 32:69-74.
- 10 42. Prabhu DC, Valechha A, Wanjari S, et al. Carbon composite beads for immobilization of
11 carbonic anhydrase. *J. Mol. Catal. B Enzym.* 2011;71:71-78.
- 12 43. Kim IG, Jo BH, Kang DG, et al. Biomineralization-based conversion of carbon dioxide to
13 calcium carbonate using recombinant carbonic anhydrase. *Chemosphere* 2012;87(10):1091-1096.
- 14 44. Jo BH, Seo JH, Yang YJ, et al. Bioinspired Silica Nanocomposite with Autoencapsulated
15 Carbonic Anhydrase as a Robust Biocatalyst for CO₂ Sequestration. *ACS Catal.*
16 2014;4(12):4332-4340.

17
18
19
20
21
22




1
2
3
4

Fig. 1. Schematic of isolation, optimization and CaCO_3 production using carbonic anhydrase.




5
6
7

Fig. 2. Flowchart summarising the overall research methodology.




15
 16
 17 **Fig. 3.** (a) Selection of bacterial isolate on the basis of CA activity. (b) Microscopic view of
 18 bacterial isolate T5.
 19



20 **Fig. 4.** (a) Gel image of 16S rDNA amplicon [Lane 1: 16S rDNA amplicon and Lane 2: DNA
 21 markers]. (b) Phylogenetic tree of bacterial T5 isolate.
 22
 23
 24

1
2




3
4

5 **Fig. 5.** (a) Effect of different media (Luria broth [LB], nutrient broth [NB], Peptone broth [PB],
6 minimal salt media [MS], basal salt media [BS], muller Hinton broth [MB]) on the production of
7 carbonic anhydrase was determined, with activity at nutrient broth as 100 %. (b) Effect of
8 incubation temperature on carbonic anhydrase production was determined, with activity at 40 °C
9 as 100 %. (c) Effect of inoculum volume (2, 4, 6, 8, 10, 12 % v/v) on CA production was
10 calculated, with activity at 4 % v/v) as 100 %. (d) Effect of agitation rate on CA production was
11 determined, with activity at 120 rpm as 100 %.

12
13
14
15
16
17

1
2



3 **Fig. 6.** (a) Effect of pH on activity. The enzyme activity at different pH (sodium citrate buffer of
4 pH 4.0-5.5, Phosphate buffer of pH 6.0-7.5 and, Tris-HCl buffer of pH 8.0-9.5) was determined,
5 with the activity at pH 7 as 100%. (b) The effect of different temperatures (25- 60°C on the
6 activity was measured at pH 7.0, with the activity at 35°C as 100%. (c) The effect of reaction
7 time on the activity was measured with the activity at 10 min as 100%. (d) Effect of organic
8 solvents on CA activity was determined.

9
10
11
12
13
14

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42





Fig. 7. SEM image of CaCO₃ precipitate (a) Spherical vaterite formed in control (b) Spherical vaterite and rhombohedral calcite crystal formed in presence of enzyme (c) XRD analyses of CaCO₃ precipitates.

1 **Table 1.** Activity of Bacterial CA in The Presence of Selected Metal Ions

2

Metal ion	Concentration (mM)	Relative Activity (%) ³
NaCl	1	87.43
	5	82.16
KCl	1	66.76
	5	70.69
MgCl ₂	1	54.45
	5	48.88
ZnCl ₂	1	103.75
	5	106.00
CaCl ₂	1	100.42
	5	99.30
FeCl ₃	1	101.64
	5	103.71
AlCl ₃	1	44.41
	5	37.47