

Electronic Supplementary Material

Regeneration of nitrate and phosphate from toilet wastewater using waste alumina adsorbent for cultivation of *Spirulina platensis*

Sang-Jun Lee^{1,2*}, Seonghwan Park^{1,2*}, Won Noh^{1,2}, Dong-Hyuk Yeom², Sooyeon Kim², Dae-wook Kim², Jungmin Kim^{1,3†}

¹Biomass Research Group, Gyeongnam Department of Environmental Toxicology & Chemistry, Korea Institute of Toxicology, 17, Jegok-gil, Samgok-ri, Munsan-eup, Jinju-si, Gyeongsangnam-do 52834, Republic of Korea

²Eco-Toxicology Research Group, Gyeongnam Department of Environmental Toxicology & Chemistry, Korea Institute of Toxicology, 17, Jegok-gil, Samgok-ri, Munsan-eup, Jinju-si, Gyeongsangnam-do 52834, Republic of Korea

³Environmental Biology Research Group, Gyeongnam Department of Environmental Toxicology & Chemistry, Korea Institute of Toxicology, 17, Jegok-gil, Samgok-ri, Munsan-eup, Jinju-si, Gyeongsangnam-do 52834, Republic of Korea

*Sang-Jun Lee and Seonghwan Park equally contributed to the first authorship.

*** Corresponding author**

Jungmin Kim, Tel: +82-55-750-3766, Fax: +82-55-750-3799

E-mail: jungmin.kim@kitox.re.kr

Figure legends

Fig. S1. Procedure used to prepare regenerated water (RW) from wastewater (WW).

Fig. S2. Microscopic and cultivation images of *Spirulina platensis*.

Fig. S3. Time course of color changes of *S. Platensis* cultures in different media.

Fig. S4. Image of *Daphnia magna* acute immobilization test.




Fig. S1. Procedure used to prepare regenerated water (RW) from wastewater (WW).




Fig. S2. Microscopic and cultivation images of *Spirulina platensis*.




Fig. S3. Time course of color changes of *S. Platensis* cultures in different media.




Fig. S4. Image of *Daphnia magna* acute immobilization test.